

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSPTAALF1626

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	DEC 01	ChemPort single article sales feature unavailable
NEWS	3	FEB 02	Simultaneous left and right truncation (SLART) added for CERAB, COMPUAB, ELCOM, and SOLIDSTATE
NEWS	4	FEB 02	GENBANK enhanced with SET PLURALS and SET SPELLING
NEWS	5	FEB 06	Patent sequence location (PSL) data added to USGENE
NEWS	6	FEB 10	COMPENDEX reloaded and enhanced
NEWS	7	FEB 11	WTEXTILES reloaded and enhanced
NEWS	8	FEB 19	New patent-examiner citations in 300,000 CA/CAPLUS patent records provide insights into related prior art
NEWS	9	FEB 19	Increase the precision of your patent queries -- use terms from the IPC Thesaurus, Version 2009.01
NEWS	10	FEB 23	Several formats for image display and print options discontinued in USPATFULL and USPAT2
NEWS	11	FEB 23	MEDLINE now offers more precise author group fields and 2009 MeSH terms
NEWS	12	FEB 23	TOXCENTER updates mirror those of MEDLINE - more precise author group fields and 2009 MeSH terms
NEWS	13	FEB 23	Three million new patent records blast AEROSPACE into STN patent clusters
NEWS	14	FEB 25	USGENE enhanced with patent family and legal status display data from INPADOCDB
NEWS	15	MAR 06	INPADOCDB and INPAFAMDB enhanced with new display formats
NEWS	16	MAR 11	EPFULL backfile enhanced with additional full-text applications and grants
NEWS	17	MAR 11	ESBIOBASE reloaded and enhanced
NEWS	18	MAR 20	CAS databases on STN enhanced with new super role for nanomaterial substances
NEWS	19	MAR 23	CA/CAPLUS enhanced with more than 250,000 patent equivalents from China
NEWS	20	MAR 30	IMSPATENTS reloaded and enhanced
NEWS	21	APR 03	CAS coverage of exemplified prophetic substances enhanced
NEWS	22	APR 07	STN is raising the limits on saved answers
NEWS	23	APR 24	CA/CAPLUS now has more comprehensive patent assignee information
NEWS	24	APR 26	USPATFULL and USPAT2 enhanced with patent assignment/reassignment information
NEWS	25	APR 28	CAS patent authority coverage expanded
NEWS	26	APR 28	ENCOMPLIT/ENCOMPLIT2 search fields enhanced
NEWS	27	APR 28	Limits doubled for structure searching in CAS REGISTRY
NEWS	28	MAY 08	STN Express, Version 8.4, now available
NEWS	29	MAY 11	STN on the Web enhanced

NEWS 30 MAY 11 BEILSTEIN substance information now available on
STN Easy

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN customer
agreement. This agreement limits use to scientific research. Use
for software development or design, implementation of commercial
gateways, or use of CAS and STN data in the building of commercial
products is prohibited and may result in loss of user privileges
and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 17:16:10 ON 13 MAY 2009

=> file capl

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.22	0.22

FILE 'CAPLUS' ENTERED AT 17:16:20 ON 13 MAY 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.
The CA Lexicon is the copyrighted intellectual property of the
American Chemical Society and is provided to assist you in searching
databases on STN. Any dissemination, distribution, copying, or storing
of this information, without the prior written consent of CAS, is
strictly prohibited.

FILE COVERS 1907 - 13 May 2009 VOL 150 ISS 20

FILE LAST UPDATED: 12 May 2009 (20090512/ED)

REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2009

USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2009

CPlus now includes complete International Patent Classification (IPC)
reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate

=> s 2006-591658/apps

0 2006-591658/AP

0 2006-591658/PRN

L1 0 2006-591658/APPS

(2006-591658/AP,PRN)

```
=> s us 20080269184/pn
L2          1 US 20080269184/PN
           (US20080269184/PN)
```

```
=> select l2
ENTER ANSWER NUMBER OR RANGE (1-):1
ENTER DISPLAY CODE (TI) OR ?:rn
E1 THROUGH E30 ASSIGNED
```

```
=> s e1-e30
      58672 112-80-1/BI
      4773 1121-60-4/BI
      6315 1338-43-8/BI
      138 143-02-2/BI
      87 143-03-3/BI
     21444 143-07-7/BI
      297 14982-53-7/BI
     1356 151-41-7/BI
     1167 313-04-2/BI
     2374 361-09-1/BI
      60 40904-90-3/BI
     137 4754-44-3/BI
     396 516-95-0/BI
     26127 544-63-8/BI
     51475 57-10-3/BI
     63354 57-11-4/BI
    141242 57-88-5/BI
      488 6156-78-1/BI
      11 65028-70-8/BI
     7061 67-97-0/BI
     5401 691397-13-4/BI
      43 71794-64-4/BI
      54 72924-08-4/BI
     766 7789-46-0/BI
     2640 80-97-7/BI
      3 823808-59-9/BI
      1 864444-61-1/BI
     3770 872-85-5/BI
    17791 9005-65-6/BI
     4769 9005-67-8/BI
L3      296106 (112-80-1/BI OR 1121-60-4/BI OR 1338-43-8/BI OR 143-02-2/BI OR
              143-03-3/BI OR 143-07-7/BI OR 14982-53-7/BI OR 151-41-7/BI OR
              313-04-2/BI OR 361-09-1/BI OR 40904-90-3/BI OR 4754-44-3/BI OR
              516-95-0/BI OR 544-63-8/BI OR 57-10-3/BI OR 57-11-4/BI OR 57-88-
              5/BI OR 6156-78-1/BI OR 65028-70-8/BI OR 67-97-0/BI OR 691397-13
              -4/BI OR 71794-64-4/BI OR 72924-08-4/BI OR 7789-46-0/BI OR 80-97
              -7/BI OR 823808-59-9/BI OR 864444-61-1/BI OR 872-85-5/BI OR 9005
              -65-6/BI OR 9005-67-8/BI)
```

```
=> FIL REGISTRY
```

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	75.52	75.74

FILE 'REGISTRY' ENTERED AT 17:18:18 ON 13 MAY 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 12 MAY 2009 HIGHEST RN 1146247-90-6
DICTIONARY FILE UPDATES: 12 MAY 2009 HIGHEST RN 1146247-90-6

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> S 14982-53-7/RN

L4 1 14982-53-7/RN

=> SET NOTICE 1 DISPLAY

NOTICE SET TO 1 U.S. DOLLAR FOR DISPLAY COMMAND
SET COMMAND COMPLETED

=> D L4 SQIDE 1-

YOU HAVE REQUESTED DATA FROM 1 ANSWERS - CONTINUE? Y/(N):y
THE ESTIMATED COST FOR THIS REQUEST IS 6.85 U.S. DOLLARS
DO YOU WANT TO CONTINUE WITH THIS REQUEST? (Y)/N:y

L4 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2009 ACS on STN

RN 14982-53-7 REGISTRY

CN Cholestane (CA INDEX NAME)

OTHER NAMES:

CN (8R,9S,10S,13R,14S,17R)-17-[(1R)-1,5-Dimethylhexyl]hexadecahydro-10,13-dimethyl-1H-cyclopenta[a]phenanthrene

CN 1H-Cyclopenta[a]phenanthrene, 17-[(1R)-1,5-dimethylhexyl]hexadecahydro-10,13-dimethyl-, (8R,9S,10S,13R,14S,17R)-

CN NSC 140722

FS STEREOSEARCH

MF C27 H48

CI COM

LC STN Files: AGRICOLA, ANABSTR, BEILSTEIN*, BIOSIS, BIOTECHNO, CA, CAPLUS, CASREACT, CHEMLIST, CIN, EMBASE, IFICDB, IFIPAT, IFIUDB, PROMT, SPECINFO, TOXCENTER, TULSA, USPAT2, USPATFULL, USPATOLD
(*File contains numerically searchable property data)

Other Sources: EINECS**

(**Enter CHEMLIST File for up-to-date regulatory information)

DT.CA CAplus document type: Conference; Dissertation; Journal; Patent; Report


RL.P Roles from patents: ANST (Analytical study); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT (Reactant or reagent); USES (Uses)

RLD.P Roles for non-specific derivatives from patents: ANST (Analytical study); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)

RL.NP Roles from non-patents: ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); MSC (Miscellaneous); OCCU

(Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT
 (Reactant or reagent); USES (Uses); NORL (No role in record)
 RLD.NP Roles for non-specific derivatives from non-patents: ANST (Analytical
 study); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU
 (Occurrence); PREP (Preparation); PROC (Process); PRP (Properties); RACT
 (Reactant or reagent); USES (Uses)

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

297 REFERENCES IN FILE CA (1907 TO DATE)
 37 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 297 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> SET NOTICE LOGIN DISPLAY

NOTICE SET TO OFF FOR DISPLAY COMMAND
 SET COMMAND COMPLETED

=>

=> s 13 and porphyrin


1 112-80-1/BI
 (112-80-1/RN)
 1 1121-60-4/BI
 (1121-60-4/RN)
 1 1338-43-8/BI
 (1338-43-8/RN)
 1 143-02-2/BI
 (143-02-2/RN)
 1 143-03-3/BI
 (143-03-3/RN)
 1 143-07-7/BI
 (143-07-7/RN)
 1 14982-53-7/BI
 (14982-53-7/RN)
 1 151-41-7/BI
 (151-41-7/RN)
 1 313-04-2/BI
 (313-04-2/RN)
 1 361-09-1/BI
 (361-09-1/RN)
 1 40904-90-3/BI
 (40904-90-3/RN)
 1 4754-44-3/BI
 (4754-44-3/RN)

1 516-95-0/BI
(516-95-0/RN)
1 544-63-8/BI
(544-63-8/RN)
1 57-10-3/BI
(57-10-3/RN)
1 57-11-4/BI
(57-11-4/RN)
1 57-88-5/BI
(57-88-5/RN)
1 6156-78-1/BI
(6156-78-1/RN)
1 65028-70-8/BI
(65028-70-8/RN)
1 67-97-0/BI
(67-97-0/RN)
1 691397-13-4/BI
(691397-13-4/RN)
1 71794-64-4/BI
(71794-64-4/RN)
1 72924-08-4/BI
(72924-08-4/RN)
1 7789-46-0/BI
(7789-46-0/RN)
1 80-97-7/BI
(80-97-7/RN)
1 823808-59-9/BI
(823808-59-9/RN)
1 864444-61-1/BI
(864444-61-1/RN)
1 872-85-5/BI
(872-85-5/RN)
1 9005-65-6/BI
(9005-65-6/RN)
1 9005-67-8/BI
(9005-67-8/RN)
5742 PORPHYRIN
3 PORPHYRINS
5742 PORPHYRIN
(PORPHYRIN OR PORPHYRINS)

L5 1 L3 AND PORPHYRIN

=> d

L5 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2009 ACS on STN
RN 40904-90-3 REGISTRY
ED Entered STN: 16 Nov 1984
CN 21H,23H-Porphine, 5,10,15,20-tetra-2-pyridinyl- (CA INDEX NAME)
OTHER NAMES:
CN 5,10,15,20-Tetra-2-pyridylporphine
CN 5,10,15,20-Tetrakis(2-pyridyl)porphyrin
CN meso-Tetra-2-pyridylporphine
CN meso-Tetrakis(2-pyridyl)porphyrin
CN meso-Tetrakis(o-pyridyl)porphine
MF C40 H26 N8
CI COM
LC STN Files: BEILSTEIN*, CA, CAPLUS, CASREACT, CHEMCATS, TOXCENTER,
USPAT2, USPATFULL
(*File contains numerically searchable property data)



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

60 REFERENCES IN FILE CA (1907 TO DATE)
 2 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA
 60 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> file capl

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

10.41

86.15

FILE 'CAPLUS' ENTERED AT 17:19:22 ON 13 MAY 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 13 May 2009 VOL 150 ISS 20

FILE LAST UPDATED: 12 May 2009 (20090512/ED)

REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2009

USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2009

Caplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate

=> s 13 and porphyrin
 40651 PORPHYRIN
 27798 PORPHYRINS
 47715 PORPHYRIN
 (PORPHYRIN OR PORPHYRINS)
 L6 758 L3 AND PORPHYRIN

=> s 13 and niosome
 359 NIOSOME
 479 NIOSOMES
 500 NIOSOME
 (NIOSOME OR NIOSOMES)
 L7 241 L3 AND NIOSOME

=> s 16 and 17
 L8 2 L6 AND L7

=> d 1-2 bib abs

L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 2005:1004559 CAPLUS
 DN 143:292573
 TI Niosome having metal porphyrin complex embedded
 therein, process for producing the same and drug with the use thereof
 IN Yuasa, Makoto; Oyaizu, Kenichi; Yamaguchi, Aritomo; Hanyuu, Yukihiro;
 Kasahara, Kazunori; Komuro, Masayasu
 PA Japan
 SO PCT Int. Appl., 40 pp.
 CODEN: PIXXD2
 DT Patent
 LA Japanese
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005084665	A1	20050915	WO 2004-JP2750	20040304
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	EP 1731150	A1	20061213	EP 2004-717289	20040304
	R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
	CN 1942184	A	20070404	CN 2004-80042914	20040304
	KR 2007008623	A	20070117	KR 2006-720709	20061002
	US 20080269184	A1	20081030	US 2007-591658	20070815
PRAI	WO 2004-JP2750	W	20040304		

OS MARPAT 143:292573

AB Disclosed is a niosome having a metal porphyrin complex embedded therein which contains a cationized metal porphyrin complex and a niosome-forming substance. This niosome having a metal porphyrin complex embedded therein has an SOD activity and can target super oxide anion radical (O2-.) and surely decrease it. Because of being in the form of a niosome, it can be delivered to, for example, a cancer cell in vivo. Therefore, it can decrease O2-. in a cancer cell and exert an

excellent effect of treating cancer. Moreover, it shows a selective effect and, therefore, is usable as a novel anticancer agent with no side effect. In addition, it can be hold in the blood, which makes it favorable as an antioxidant. Owing to this characteristic, it can protect the living body from in vivo disorders caused by active oxygen. For example, iron[5,10,15,20-tetrakis(2-methylpyridyl)porphyrin] was prepared, and mixed with stearic acid metal salt to form an ion complex of the porphyrin. Then, the ion complex was mixed with tween-61 and cholesterol to form a niosome to exam for its antitumor activity and antioxidant activity in vitro.

RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2009 ACS on STN
AN 2005:674689 CAPLUS
DN 144:156398
TI Novel functional nano-size nonionic surfactant particles on which cationic metalloporphyrins are absorbed; preparation, characterization, and antioxidant properties
AU Yuasa, Makoto; Oyaizu, Kenichi; Hanyuu, Yukihiro; Kasahara, Kazunori; Yamaguchi, Aritomo
CS Department of Pure and Applied Chemistry, Faculty of Science and Technology, Tokyo University of Science, Noda, 278-8510, Japan
SO Journal of Oleo Science (2005), 54(8), 465-471
CODEN: JOSOAP; ISSN: 1345-8957
PB Japan Oil Chemists' Society
DT Journal
LA Japanese
AB Cationic manganese(III) 5,10,15,20-tetrakis(N-methylpyridinium-2-yl) porphyrin (MnT2MPyP) and manganese(III) 5,10,15,20-tetrakis(N-methylpyridinium-4-yl) porphyrin (MnT4MPyP) complexes were synthesized as superoxide dismutase (SOD) mimics which were introduced into niosomes to examine this effects on the capacity of drug delivery system (DDS) to maintain and perpetuate blood circulation. All the niosomes were prepared from polyoxyethylene sorbitan monostearate (Tween 61) by the conventional sonication method. SOD activity was measured by the stopped-flow anal. and the cytochrome c method. Sodium stearate-linked MnT2MPyP was the most effective catalyst along with SOD activity for decomposing O₂⁻ · at a second-order rate constant of 2.0×10⁷ M⁻¹ s⁻¹ in Tween 61 niosomes. Rate consts. of metalloporphyrin-embedded niosomes for reaction with hydrogen peroxide (H₂O₂) and half-life times in H₂O₂ were also determined Metalloporphyrin-embedded niosomes were found to have greater half-life times compared to metalloporphyrin without niosomes. The present findings clearly indicate that metalloporphyrin-embedded niosomes are highly effective for bringing about O₂⁻ · decomposition and should thus find application as DDS in antioxidant drugs.

=> s 13 and ((liposome OR "Pharmaceutical liposomes") OR "Liposomes")
40348 LIPOSOME
52181 LIPOSOMES
59952 LIPOSOME
(LIPOSOME OR LIPOSOMES)
341449 "PHARMACEUTICAL"
93542 "PHARMACEUTICALS"
397390 "PHARMACEUTICAL"
("PHARMACEUTICAL" OR "PHARMACEUTICALS")
52181 "LIPOSOMES"
4949 "PHARMACEUTICAL LIPOSOMES"
("PHARMACEUTICAL" (W) "LIPOSOMES")

52181 "LIPOSOMES"
L9 10368 L3 AND ((LIPOSOME OR "PHARMACEUTICAL LIPOSOMES") OR "LIPOSOMES")

=> s 16 and 19

L10 59 L6 AND L9

=> s l10 and drug delivery

864061 DRUG
380271 DRUGS
1044341 DRUG
(DRUG OR DRUGS)
330079 DELIVERY
2027 DELIVERIES
331301 DELIVERY
(DELIVERY OR DELIVERIES)
233444 DRUG DELIVERY
(DRUG(W)DELIVERY)

L11 19 L10 AND DRUG DELIVERY

=> d 1-19 bib hitstr abs

L11 ANSWER 1 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2007:350750 CAPLUS

DN 147:307839

TI Design of pH-sensitive nano-carrier with control ability of in vivo
free-radicals

AU Kawakami, Hiroyoshi

CS Department of Applied Chemistry, Tokyo Metropolitan University, Hachioji,
Tokyo, 192-0397, Japan

SO Maku (2006), 31(6), 290-295

CODEN: MAKUD9; ISSN: 0385-1036

PB Nippon Maku Gakkai

DT Journal

LA Japanese

IT 9005-65-6, Tween-80.

RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL
(Biological study); USES (Uses)

(pH-sensitive liposomes with control ability of in vivo
free-radicals in the treatment of cancer)

RN 9005-65-6 CAPLUS

CN Sorbitan, mono-(9Z)-9-octadecenoate, poly(oxy-1,2-ethanediyl) derivs. (CA
INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

AB Liposomes are well known to be drug carriers of the phospholipid
bilayer with potential application in drug and gene deliveries. However,
the cellular uptake of liposomes generally follows an endocytic
pathway so that most liposomes remain entrapped endosomes and
are unable to reach the cytoplasmic space. To overcome this problem,
pH-sensitive liposomes, which are designed to undergo rapid
destabilization in the acidic environments of endosomes, have been
synthesized. We report the novel design of an antioxidant or anticancer
drug delivery system based on a pH-sensitive
liposome retaining the metalloporphyrin as an SOD mimic. The
liposomes contained cationic/anionic lipid combinations and were
composed of Fe-porphyrin, L- α -phosphatidylcholine (DMPC),
dimethylditetradecylammonium bromide (DTDAB), sodium oleate (OANa), and
Tween-80. The size of the liposome was approx. 30 nm,
indicating that the resulting liposome was a nanoparticle.

L11 ANSWER 2 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2007:155681 CAPLUS

DN 146:407827
TI pH-sensitive liposome retaining Fe-porphyrin as SOD
mimic for novel anticancer drug delivery system
AU Kawakami, Hiroyoshi; Hiraka, Kazue; Tamai, Miho; Horiuchi, Aiko; Ogata,
Akihiko; Hatsugai, Tomomi; Yamaguchi, Aritomo; Oyaizu, Kenichi; Yuasa,
Makoto
CS Department of Applied Chemistry, Tokyo Metropolitan University, Hachioji,
Tokyo, 192-0397, Japan
SO Polymers for Advanced Technologies (2007), 18(1), 82-87
CODEN: PADTE5; ISSN: 1042-7147
PB John Wiley & Sons Ltd.
DT Journal
LA English
IT 9005-65-6, Tween-80
RL: PEP (Physical, engineering or chemical process); PRP (Properties); THU
(Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(pH-sensitive liposome retaining iron-porphyrin as
SOD mimic for anticancer drug delivery system)
RN 9005-65-6 CAPLUS
CN Sorbitan, mono-(9Z)-9-octadecenoate, poly(oxy-1,2-ethanediyl) derivs. (CA
INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

AB In this article the novel design of an anticancer drug
delivery system is reported based on a pH-sensitive
liposome retaining the Fe-porphyrin as a superoxide
dismutase(SOD) mimic. The liposomes contained cationic/anionic
lipid combinations and were composed of Fe-porphyrin,
L- α -phosphatidylcholine, dimethylditetradecylammonium bromide,
sodium oleate, and Tween-80. The size of the liposome was
approx. 30 nm. The EC50 value (the effective concentration of compound
required to
produce a 50% LD against cells) of the liposome was found to be
significantly smaller than that of cisplatin as the control drug,
suggesting that the liposome showed a high cytotoxicity toward
the cancer cells. This is due to the fact that the pH-sensitive
liposome rapidly corresponds to the acidic environments of the
endosomes and is unstable, and the Fe-porphyrin is delivered
into the cytosol. This results suggests that O₂⁻ may be useful as a
target mol. to induce the selective death of cancer cells and that a
pH-sensitive liposome retaining Fe-porphyrin as an SOD
mimic is a new class of anticancer agent.

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
AN 2007:133743 CAPLUS
DN 148:17232
TI Synthesis of cationic manganese porphyrin bearing alkylsulfonio
groups and evaluation of their antioxidant activities
AU Yuasa, Makoto; Oyaizu, Kenichi; Murata, Hidenori; Komuro, Masayasu; Awa,
Ryota; Ohkubo, Ayumi
CS Department of Pure and Applied Chemistry, Faculty of Science and
Technology, Tokyo University of Science, Noda, 278-8510, Japan
SO Journal of Oleo Science (2007), 56(2), 95-101
CODEN: JOSOAP; ISSN: 1345-8957
PB Japan Oil Chemists' Society
DT Journal
LA Japanese
IT 691397-13-4, Pluronic F-68
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL
(Biological study); USES (Uses)

(synthesis of cationic manganese porphyrin bearing
alkylsulfonio groups and evaluation of their antioxidant activities)


RN 691397-13-4 CAPLUS

CN Oxirane, 2-methyl-, polymer with oxirane, triblock (CA INDEX NAME)

CM 1

CRN 75-56-9

CMF C3 H6 O



CM 2

CRN 75-21-8

CMF C2 H4 O



AB A water-soluble cationic 5,10,15,20-tetrakis(2-dimethylsulfoniophenyl)-porphinatomanganese (III) ion (MnT2M2SuP) and a 5,10,15,20-tetrakis(4-dimethylsulfoniophenyl)porphinatomanganese(III) ion (MnT4M2SuP) were synthesized as superoxide dismutase (SOD) mimics which were introduced into PEG- liposome composed of dimyristoylphosphatidylcholine (DMPC) and Pluronic F-68 to examine the effect of the liposome on the capacity for use as drug delivery system (DDS) to maintain and perpetuate blood circulation. Fluorescence spectra in pseudo blood circulation expts. indicated that MnT4M2SuP continued to be bundled in PEG-liposome, while fluorescence from cross-section of cell observed by confocal laser scanning microscope indicated that PEG-liposome was ingested into a cell. SOD activity was determined by stopped-flow anal., which allowed the determination of kcat values for the reaction of the metalloporphyrins with superoxide anion radical ($\cdot\text{O}_2^-$). Solution of PEG-liposome loaded with MnT2M2SuP or MnT4M2SuP were the most effective catalyst as a SOD mimic to decompose $\cdot\text{O}_2^-$ at second-order rate consts. of 3.5-4.5 + 107 M⁻¹ s⁻¹.

L11 ANSWER 4 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2007:133674 CAPLUS

DN 147:371446

TI Antioxidant and anticancer properties of metalloporphyrins embedded in liposomes

AU Yuasa, Makoto; Oyaizu, Kenichi; Murata, Hidenori; Sahara, Yoshizumi; Hatsugai, Tomomi; Ogata, Akihiko

CS Department of Pure and Applied Chemistry, Faculty of Science and Technology, Tokyo University of Science, Noda, 278-8510, Japan

SO Journal of Oleo Science (2007), 56(2), 87-93

CODEN: JOSOAP; ISSN: 1345-8957

PB Japan Oil Chemists' Society

DT Journal

LA Japanese


IT 65028-70-8

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);

PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(antioxidant and anticancer properties of metalloporphyrins embedded in liposomes)

RN 65028-70-8 CAPLUS

CN Manganese(4+), [[2,2',2'',2'''-(21H,23H-porphine-5,10,15,20-tetrayl)tetrakis[1-methylpyridiniumato]](2-)-κN21,κN22,κN23,κN24]-, (SP-4-1)- (9CI) (CA INDEX NAME)



IT 691397-13-4, Pluronic F-68

RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(antioxidant and anticancer properties of metalloporphyrins embedded in liposomes)


RN 691397-13-4 CAPLUS

CN Oxirane, 2-methyl-, polymer with oxirane, triblock (CA INDEX NAME)

CM 1

CRN 75-56-9

CMF C3 H6 O



CM 2

CRN 75-21-8

CMF C2 H4 O



AB Reactive oxygen species (ROS) are implicated in many disease such as

inflammation, arteriosclerosis, cancer. Therefore, a water-soluble cationic metalloporphyrins with SOD activity are studied widely as antioxidant drugs. Further, liposomes are applied to drug delivery system (DDS) as drug carriers and investigated for example disposition and stability. We designed PEG modified liposomes for avoiding reticuloendothelial system (RES) and embedded cationic metalloporphyrins for DDS, evaluated antioxidant and anticancer property. Preservation of these particle size measured DLS in an in vitro system, in order to simulate in vivo conditions of flow. Result of this measurement, we found Pluronic F-68/ liposomes have a long circulation property, and avoid fusion with plasma protein. SOD activity was determined by the stopped-flow anal. and cytochrome c assay, which allowed the evaluation of kcat and IC50 for the reaction with a superoxide anion radical ($\cdot\text{O}_2^-$). Anti cancer property was measured by cell viability test. We found that F-68/ liposomes were the most effective catalyst as antioxidant and anticancer. These results revealed that porphyrin-embedded PEG-liposomes had the property of long circulation in blood and that this compound was effective as a SOD model compound with a drug carrier capacity.

L11 ANSWER 5 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 2006:379466 CAPLUS
 DN 145:460088
 TI Pegylated tetraarylporphyrin entrapped in liposomal membranes
 AU Kepczynski, Mariusz; Nawalany, Kinga; Jachimska, Barbara; Romek, Marek; Nowakowska, Maria
 CS Faculty of Chemistry, Jagiellonian University, Krakow, 30-060, Pol.
 SO Colloids and Surfaces, B: Biointerfaces (2006), 49(1), 22-30
 CODEN: CSBBEQ; ISSN: 0927-7765
 PB Elsevier B.V.
 DT Journal
 LA English
 IT 57-10-3, Palmitic acid, biological studies 57-11-4, Stearic acid, biological studies 112-80-1, Oleic acid, biological studies
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (pegylated tetraarylporphyrin entrapped in liposomal membranes for photodynamic therapy)
 RN 57-10-3 CAPLUS
 CN Hexadecanoic acid (CA INDEX NAME)


$\text{HO}_2\text{C}-(\text{CH}_2)_{14}-\text{Me}$

RN 57-11-4 CAPLUS
 CN Octadecanoic acid (CA INDEX NAME)

$\text{HO}_2\text{C}-(\text{CH}_2)_{16}-\text{Me}$

RN 112-80-1 CAPLUS
 CN 9-Octadecenoic acid (9Z)- (CA INDEX NAME)

Double bond geometry as shown.



AB A system of poly(ethylene glycol) bound tetraarylporphyrin entrapped in liposomal membranes was investigated. The interactions between the 5-(4-hydroxymethylphenyl)-10,15,20-tritolylporphyrin (Po) covalently attached to the poly(ethylene glycol) chain (PEG-Po), and phosphatidylcholine liposomes in the aqueous solution were studied. The adsorption of the investigated polymer to lipid vesicles was confirmed by measurements of dynamic light scattering and zeta potential. Exptl. results demonstrate that the diameter of liposomes increased and the absolute value of the zeta potential decreased after addition of PEG-Po.

The

binding consts. (Kb) of Po chromophores to liposome in pH range from 5.2 to 9.0 were determined using fluorescence spectroscopy. The degree of binding was found to be pH-independent and the average value was 24.6 ± 0.9 mg ml⁻¹. The acid-base properties of the porphyrin chromophores and their aggregation in an aqueous solution were also studied. pK values associated

with imine-N protonation of the porphyrin core were found to be 2.59 and 0.68 at the ionic strength of 0.1 M. The equilibrium constant for dimerization, KD, was found to be 5×10^3 M⁻¹.

RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 6 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2006:210135 CAPLUS

DN 144:260853

TI Liposome compositions for cancer therapy, and manufacture thereof

IN Aoki, Yoichi; Ueda, Eiichi

PA Konica Minolta Medical & Graphic, Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 18 pp.

CODEN: JKXXAF


DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2006063009	A	20060309	JP 2004-246737	20040826
PRAI	JP 2004-246737		20040826		
IT	57-88-5, Cholest-5-en-3-ol (3 β)-, biological studies				
	RL: PEP (Physical, engineering or chemical process); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)				
	(liposome comps. for cancer therapy, and manufacture thereof)				
RN	57-88-5 CAPLUS				
CN	Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)				

Absolute stereochemistry.



AB The invention relates to a liposome for cancer therapy, formed from polyethylene glycol group-containing compound with other phospholipid membrane components by using supercrit. or subcrit. carbon dioxide, wherein the liposome has an average particle size 0.75-0.85 μm and contains an active component in the lipid membrane or inner water phase with no organic solvent. The liposome shows improved stability of the active component, and enables efficient delivery of the active component. A method for manufacturing the liposome is also disclosed. For example, a liposome was prepared from dipalmitoylphosphatidylcholine, cholesterol, ethoxylated phospholipid (Sunbright DSPE-020CN), and adriamycin by using supercrit. carbon dioxide.

L11 ANSWER 7 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2006:210040 CAPLUS

DN 144:260852

TI Liposome compositions for cancer therapy, and manufacture thereof

IN Aoki, Yoichi; Ueda, Eiichi

PA Konica Minolta Medical & Graphic, Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 18 pp.

CODEN: JKXXAF


DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2006063008	A	20060309	JP 2004-246736	20040826
PRAI	JP 2004-246736		20040826		
IT	57-88-5, Cholesterol, biological studies				
	RL: PEP (Physical, engineering or chemical process); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)				
	(liposome compns. for cancer therapy, and manufacture thereof)				
RN	57-88-5 CAPLUS				
CN	Cholest-5-en-3-ol (3β)- (CA INDEX NAME)				


Absolute stereochemistry.




AB The invention relates to a liposome having an average particle size 0.75-0.85 μm and containing an active component in the lipid membrane or inner water phase with no organic solvent. The liposome shows improved stability of the active component, and is suitable for use for intraarterial injection chemotherapy, neutron capture therapy, and phototherapy. A method for manufacturing the liposome by using supercrit. or subcrit. carbon dioxide is also disclosed. For example, a liposome was prepared from dipalmitoylphosphatidylcholine, cholesterol, ethoxylated phospholipid (Sunbright DSPE-020CN), and iopamidol by using supercrit. carbon dioxide.

L11 ANSWER 8 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 2005:641688 CAPLUS
 DN 143:208263
 TI Topical use of liposomal copper palmitate formulation blocks
 porphyrin-induced photosensitivity in rats
 AU Bilgin, Mehmet D.; Elcin, A. Eser; Elcin, Y. Murat
 CS Department of Biophysics, Medical Faculty, Adnan Menderes University,
 Aydin, 09100, Turk.
 SO Journal of Photochemistry and Photobiology, B: Biology (2005), 80(2),
 107-114
 CODEN: JPPBEG; ISSN: 1011-1344
 PB Elsevier B.V.
 DT Journal
 LA English
 IT 57-88-5, Cholesterol, biological studies
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (as liposome component; topical liposomal copper palmitate
 blocks PDT photosensitizer-induced PpIX photosensitivity)
 RN 57-88-5 CAPLUS
 CN Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)

Absolute stereochemistry.



IT 57-10-3, Palmitic acid, reactions
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (topical liposomal copper palmitate blocks PDT photosensitizer-induced
 PpIX photosensitivity)
 RN 57-10-3 CAPLUS
 CN Hexadecanoic acid (CA INDEX NAME)



AB Photodynamic therapy (PDT) is a new treatment modality that uses
 porphyrin derivs. and visible light, especially for the treatment of
 cancer. However, PDT with certain photosensitizers can cause prolonged
 skin photosensitization. This is particularly true for Photofrin II
 (Photofrin)-mediated PDT where patients are required to avoid direct
 exposure to sunlight for a period of 4-6 wk. This is the only long-term
 adverse reaction to the drug. Recent studies have shown that topical
 copper treatment avoids this type of inflammatory reaction. In this
 study, we have tested the efficiency of the liposomal formulation of
 copper palmitate on porphyrin-photosensitized rats. Initially,
 adult male Sprague-Dawley rats were rendered photosensitive either by
 administration of Photofrin or aminolevulinic acid (ALA), a precursor of
 protoporphyrin IX (PpIX). Prior to this, their dorsal skin was shaved and
 treated topically with a cream consisting of either empty or copper
 palmitate-encapsulated liposomal formulation. After being kept in a

dimmed light environment, the rats were exposed to visible light, and inflammatory responses were inspected. Histol. studies revealed that no inflammatory cells were present at the skin sites treated with liposomal cream containing copper palmitate in the Photofrin-sensitized group while no reduction in the number of inflammatory cells was observed at the skin samples treated with the empty liposomes. In conclusion, the data demonstrate the significant protective effect of topically-applied liposome-encapsulated copper palmitate against both Photofrin and ALA-induced PpIX photosensitivity.

RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 9 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2005:300287 CAPLUS

DN 142:372464

TI Compositions comprising immunostimulatory nucleic acid-lipophilic conjugates and antigen or therapeutic agent for treating allergy, cancer, infection and autoimmune disease

IN Vollmer, Joerg; Krieg, Arthur M.; Uhlmann, Eugen

PA Coley Pharmaceutical Group, Inc., USA; Coley Pharmaceutical GmbH

SO PCT Int. Appl., 120 pp.

CODEN: PIXXD2


DT Patent

LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005030259	A2	20050407	WO 2004-US31748	20040927
	WO 2005030259	A3	20051110		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2004275876	A1	20050407	AU 2004-275876	20040927
	CA 2536139	A1	20050407	CA 2004-2536139	20040927
	US 20050130911	A1	20050616	US 2004-952254	20040927
	EP 1663316	A2	20060607	EP 2004-789138	20040927
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	JP 2007506790	T	20070322	JP 2006-528305	20040927
PRAI	US 2003-505977P	P	20030925		
	WO 2004-US31748	W	20040927		
IT	57-88-5D, Cholesterol, derivs. and conjugates 14982-53-7D, Cholestane, derivs. and conjugates				
	RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(compsns. comprising immunostimulatory nucleic acid-lipophilic conjugates and antigen or therapeutic agent for treating allergy, cancer, infection and autoimmune disease)				
RN	57-88-5 CAPLUS				
CN	Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)				

Absolute stereochemistry.



RN 14982-53-7 CAPLUS
 CN Cholestane (CA INDEX NAME)

Absolute stereochemistry.



AB The invention relates to a nucleic acid-lipophilic conjugates and methods for modulating an immune response using the conjugates. The lipophilic moiety associated with an immunostimulatory nucleic acid.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 10 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2005:34446 CAPLUS

DN 142:141238

TI Metal porphyrin complex-embedded liposomes for pharmaceuticals

IN Yuasa, Makoto; Matsukura, Noriyoshi; Yamaguchi, Aritomo; Kawakami, Hiroyoshi; Nagaoka, Shoji; Abe, Masahiko; Takebayashi, Hitoshi; Horiuchi, Aiko; Ogata, Akihiko; Sakaya, Takeshi

PA Makoto Yuasa, Japan

SO U.S. Pat. Appl. Publ., 20 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20050008687	A1	20050113	US 2004-788263	20040301
	JP 2005041869	A	20050217	JP 2004-200163	20040707
PRAI	JP 2003-193138	A	20030707		
	JP 2003-193139	A	20030707		
OS	MARPAT 142:141238				
IT	72924-08-4P				

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES


(Uses)

(metal porphyrin complex-embedded liposomes for
pharmaceuticals)

RN 72924-08-4 CAPLUS

CN Manganese(4+), [[4,4',4'',4'''-(21H,23H-porphine-5,10,15,20-tetrayl-
κN21,κN22,κN23,κN24)tetrakis[1-
methylpyridiniumato]](2-)]-, (SP-4-1)- (CA INDEX NAME)

PAGE 1-A




PAGE 2-A



IT 872-85-5, 4-Pyridylcarboxaldehyde 1121-60-4,
2-Pyridylcarboxaldehyde 6156-78-1, Manganese acetate
tetrahydrate 7789-46-0, Iron bromide (FeBr₂)
RL: RCT (Reactant); RACT (Reactant or reagent)
(metal porphyrin complex-embedded liposomes for
pharmaceuticals)


RN 872-85-5 CAPLUS

CN 4-Pyridinecarboxaldehyde (CA INDEX NAME)




RN 1121-60-4 CAPLUS

CN 2-Pyridinecarboxaldehyde (CA INDEX NAME)



RN 6156-78-1 CAPLUS
 CN Acetic acid, manganese(2+) salt, tetrahydrate (8CI, 9CI) (CA INDEX NAME)




●1/2 Mn(II)

●2 H₂O

RN 7789-46-0 CAPLUS
 CN Iron bromide (FeBr₂) (CA INDEX NAME)

Br-Fe-Br

IT 40904-90-3P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (metal porphyrin complex-embedded liposomes for
 pharmaceuticals)
 RN 40904-90-3 CAPLUS
 CN 21H,23H-Porphine, 5,10,15,20-tetra-2-pyridinyl- (CA INDEX NAME)



IT 57-10-3D, Palmitic acid, alkali metal salts 57-11-4D,
 Stearic acid, alkali metal salts 57-88-5, Cholesterol,
 biological studies 112-80-1, Oleic acid, biological studies

112-80-1D, Oleic acid, alkali metal salts 143-02-2D,
 alkali metal salts 143-03-3D, alkali metal salts
 143-07-7D, Lauric acid, alkali metal salts 151-41-7D,
 Dodecylsulfuric acid, alkali metal salts 544-63-8D, Myristic
 acid, alkali metal salts 4754-44-3D, Tetradecylsulfuric acid,
 alkali metal salts 9005-65-6, Tween 80 9005-67-8,
 Tween 61 823808-59-9D, metal complexes
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (metal porphyrin complex-embedded liposomes for
 pharmaceuticals)

RN 57-10-3 CAPLUS
 CN Hexadecanoic acid (CA INDEX NAME)




RN 57-11-4 CAPLUS
 CN Octadecanoic acid (CA INDEX NAME)




RN 57-88-5 CAPLUS
 CN Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)

Absolute stereochemistry.




RN 112-80-1 CAPLUS
 CN 9-Octadecenoic acid (9Z)- (CA INDEX NAME)

Double bond geometry as shown.



RN 112-80-1 CAPLUS
 CN 9-Octadecenoic acid (9Z)- (CA INDEX NAME)

Double bond geometry as shown.



RN 143-02-2 CAPLUS
CN 1-Hexadecanol, 1-(hydrogen sulfate) (CA INDEX NAME)

$\text{HO}_3\text{SO}-(\text{CH}_2)_{15}-\text{Me}$

RN 143-03-3 CAPLUS
CN Sulfuric acid, monooctadecyl ester (CA INDEX NAME)

$\text{HO}_3\text{SO}-(\text{CH}_2)_{17}-\text{Me}$

RN 143-07-7 CAPLUS
CN Dodecanoic acid (CA INDEX NAME)

$\text{HO}_2\text{C}-(\text{CH}_2)_{10}-\text{Me}$

RN 151-41-7 CAPLUS
CN Sulfuric acid, monododecyl ester (CA INDEX NAME)

$\text{HO}_3\text{SO}-(\text{CH}_2)_{11}-\text{Me}$

RN 544-63-8 CAPLUS
CN Tetradecanoic acid (CA INDEX NAME)

$\text{HO}_2\text{C}-(\text{CH}_2)_{12}-\text{Me}$

RN 4754-44-3 CAPLUS
CN 1-Tetradecanol, 1-(hydrogen sulfate) (CA INDEX NAME)

$\text{HO}_3\text{SO}-(\text{CH}_2)_{13}-\text{Me}$


RN 9005-65-6 CAPLUS
CN Sorbitan, mono-(9Z)-9-octadecenoate, poly(oxy-1,2-ethanediyl) derivs. (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 9005-67-8 CAPLUS
CN Sorbitan, monooctadecanoate, poly(oxy-1,2-ethanediyl) derivs. (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 823808-59-9 CAPLUS
CN Pyridinium, 4,4'-[(7,12-diethenyl-3,8,13,17-tetramethyl-21H,23H-porphine-2,18-diyl)bis[(1-oxo-3,1-propanediyl)imino]]bis[1-methyl- (9CI) (CA INDEX NAME)



AB A metalloporphyrin-complex-embedded liposome comprising a cationic metalloporphyrin complex and a lipid having liposome-forming ability is disclosed. As metalloporphyrin-complex-embedded liposomes act on superoxide anion radicals (O₂⁻), and can surely lower their concentration, they can exhibit superb effects for the treatment of cancers and have excellent characteristics as antioxidants. Thus, iron[5,10,15,20-tetrakis(2-methylpyridyl)porphyrin] was prepared starting from 2-pyridylcarboxaldehyde and pyrrole followed by reaction with FeBr₂ of the resulting porphyrin and methylation. Liposomes were obtained from the above complex and stearic acid.

L11 ANSWER 11 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 2004:837282 CAPLUS

DN 141:337627

TI Pharmaceutical composition containing artificial oxygen carrier

IN Kai, Toshiya; Katayama, Naohisa; Azuma, Yuko; Yokoe, Junichi; Kida, Yoshinori; Hukutomi, Ippei; Sato, Makoto; Tsuchida, Eishun; Takeoka, Shinji; Komatsu, Teruyuki; Sakai, Hiromi; So, Keitaro

PA Nipro Corporation, Japan

SO Eur. Pat. Appl., 21 pp.

CODEN: EPXXDW


DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1466649	A1	20041013	EP 2004-8419	20040407
	EP 1466649	B1	20080813		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
	JP 2004307404	A	20041104	JP 2003-103875	20030408
	US 20040258745	A1	20041223	US 2004-819352	20040407
	US 7417118	B2	20080826		
	AT 404245	T	20080815	AT 2004-8419	20040407
	ES 2308061	T3	20081201	ES 2004-8419	20040407
PRAI	JP 2003-103875	A	20030408		
IT	57-88-5, Cholesterol, biological studies				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(compns. containing Hb-encapsulated liposome or porphyrin-iron conjugate with albumin as oxygen carriers)				
RN	57-88-5 CAPLUS				
CN	Cholest-5-en-3-ol (3β)- (CA INDEX NAME)				

Absolute stereochemistry.



AB A pharmaceutical composition is described, containing an artificial oxygen carrier, which has high oxygen carrying efficiency; and has a required colloid osmotic pressure, an appropriate crystalloid osmotic pressure, pH, and electrolyte balance. The composition is prepared by appropriately adding at least one substance selected from the group consisting of a compound capable of providing a colloid osmotic pressure, an electrolyte, a saccharide, an amino acid, an antioxidant, a pH adjuster, and an isotonicizing agent to a preparation including a Hb-encapsulated liposome in which a Hb is encapsulated in a liposome or to a preparation including a conjugate of a porphyrin-iron complex and albumin. For example, homocysteine and an equal molar amount (relative to Hb) of a pyridoxal 5-phosphate solution were added to a Hb solution (40 g/dL), the mixture was adjusted to pH 7.4, stirred at 4° overnight, filtered, and treated with CO to yield a carbon-monoxidized Hb solution. A mixture of dipalmitoylphosphatidylcholine (DPPC)/cholesterol/dipalmitoylphosphatidylglycerol (DPPG) (10:10:2) was dissolved in benzene was freeze-dried to yield lipid-mixed powder. The lipid-mixed powder was hydrated by adding the powder to the carbon-monoxidized Hb solution and extruded to yield an HbV dispersion. Four-fold amount of physiol. saline was added to the HbV dispersion and ultrafiltrated to remove non-encapsulated Hb. The lipid concentration was adjusted to 2.0 g/dL with physiol. saline. A conjugate was formed by binding PEG and distearoylphosphatidylethanolamine via succinic acid (PEG-DSPE), dissolved in physiol. saline so as to have a PEG-DSPE content of 0.3 mol% relative to the total lipid amount, and added dropwise by portions to the HbV dispersion. The mixture was carbon monoxidized and concentrated to yield an HbV pellet layer in the lower layer. Stirred and mixed under a CO atmospheric for 2 h (37°, 400 rpm), further stirred and mixed overnight at 4°. PBS was added to the HbV pellet layer, and ultracentrifugation was performed for washing. Then, a rHSA solution (5% in PBS) was added to the lower layer to yield 200 mL of HbV solution (Hb 10 g/dL, about pH 7.4). The resultant solution had a colloid osmotic pressure of about 20 mmHg and a crystalloid osmotic pressure of about 300 mOsm.

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L11 ANSWER 12 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
AN 2003:912557 CAPLUS
DN 139:399862
TI Paramagnetic particles that provide improved relaxivity for MRI contrast agents
IN Lanza, Gregory M.; Wickline, Samuel A.
PA Barnes-Jewish Hospital, USA
SO U.S. Pat. Appl. Publ., 17 pp., Cont.-in-part of U.S. Ser. No. 153,395.
CODEN: USXXCO
DT Patent

LA English

FAN.CNT 2


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20030215392	A1	20031120	US 2003-400379	20030326
	US 7235227	B2	20070626		
	US 20030185760	A1	20031002	US 2002-153395	20020521
	US 6869591	B2	20050322		
PRAI	US 2002-368100P	P	20020326		
	US 2002-153395	A2	20020521		
IT	57-88-5, Cholesterol, biological studies 112-80-1D, Oleic acid, conjugates with gadolinium-DTPA complexes RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (paramagnetic particles that provide improved relaxivity for MRI contrast agents)				
RN	57-88-5 CAPLUS				
CN	Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)				

Absolute stereochemistry.



RN 112-80-1 CAPLUS
CN 9-Octadecenoic acid (9Z)- (CA INDEX NAME)

Double bond geometry as shown.



AB An improved contrast agent for magnetic resonance imaging comprises particles to each of which is coupled a multiplicity of chelating agents containing paramagnetic ions. In the improved agent, the position of the ion is offset from the surface of the particle so as to improve the relaxivity imparted by the contrast agent. A tether offsetting the chelate from the surface of the particle may optionally contain cleavage sites permitting more facile excretion of the chelated paramagnetic ion. Gadolinium diethylenetriaminepentaacetic acid-bis-oleate (Gd-DTPA-BOA) or DTPA-phosphatidylethanolamine (DTPA-PE), was included in the surfactant comixt. at 20 mol% of the total lipid membrane. Gadolinium chloride was added in excess proportions as a post-emulsification step to nanoparticles formulated with DTPA-PE. Unbound gadolinium was removed by dialysis on the nanoparticles against distilled deionized water (300,000 MW). Gadolinium-DTPA-BOA was incorporated into the surfactant lipids as the complete paramagnetic compound. Both Gd-DTPA-BOA and Gd-DTPA-PE emulsions were tested for free Gd³⁺ by using the arsenazo III reaction and showed no sign of unbound lanthanide. Each lipophilic nanoparticle presented more than 50,000 Gd-complexes along the water-lipid interface.


RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 13 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 2003:609843 CAPLUS
 DN 139:169326
 TI Device and methods for initiating chemical reactions and for the targeted delivery of drugs or other agents
 IN Ueberle, Friedrich
 PA Germany
 SO U.S. Pat. Appl. Publ., 19 pp.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 20030147812	A1	20030807	US 2002-316273	20021211
	EP 1319423	A2	20030618	EP 2002-27643	20021211
	EP 1319423	A3	20031008		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
PRAI	US 2001-339285P	P	20011211		
IT	67-97-0, Cholecalciferol				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (device and methods for initiating chemical reactions and for targeted delivery of drugs or other agents)				
RN	67-97-0 CAPLUS				
CN	Cyclohexanol, 3-[(2E)-2-[(1R,3aS,7aR)-1-[(1R)-1,5-dimethylhexyl]octahydro-7a-methyl-4H-inden-4-ylidene]ethylidene]-4-methylene-, (1S,3Z)- (CA INDEX NAME)				

Absolute stereochemistry.
 Double bond geometry as shown.




AB The present invention is directed to methods and apparatus for the targeted initiation or deactivation of chemical reactions by an acoustic energy source in a host. Methods and apparatus for the targeted delivery of drugs, diagnostic agents and other compds. using an acoustic energy source are also provided.

L11 ANSWER 14 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 2000:553450 CAPLUS
 DN 133:182966
 TI Novel methods of imaging and treatment with targeted compositions
 IN Ungr, Evan C.; Wu, Yunqiu

PA ImaRx Pharmaceutical Corp., USA
 SO PCT Int. Appl., 211 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 8

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000045856	A2	20000810	WO 2000-US2620	20000202
	WO 2000045856	A3	20010215		
	W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW			
	RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	US 6521211	B1	20030218	US 1999-243640	19990203
	CA 2362200	A1	20000810	CA 2000-2362200	20000202
	EP 1146911	A2	20011024	EP 2000-914480	20000202
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
	AU 777304	B2	20041007	AU 2000-35866	20000202
	AU 2005200059	A1	20050203	AU 2005-200059	20050107
PRAI	US 1999-243640	A	19990203		
	US 1995-497684	B2	19950607		
	US 1996-640464	B2	19960501		
	US 1996-660032	B2	19960606		
	US 1998-73913P	P	19980206		
	US 1998-218660	A2	19981222		
	WO 2000-US2620	W	20000202		
IT	57-88-5, Cholesterol, biological studies				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(ultrasound imaging and treatment with targeted compns.)				
RN	57-88-5 CAPLUS				
CN	Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)				

Absolute stereochemistry.




AB Novel ultrasound methods comprising administering to a patient a targeted vesicle composition which comprises vesicles comprising a lipid, protein or polymer, encapsulating a gas, in combination with a targeting ligand, and scanning the patient using ultrasound. The scanning may comprise exposing the patient to a first type of ultrasound energy and then interrogating the patient using a second type of ultrasound energy. The targeting ligand preferably targets tissues, cells or receptors, including myocardial cells, endothelial cells, epithelial cells, tumor cells and the glycoprotein GPIIbIIIa receptor. The methods may be used to detect a

thrombus, enhancement of an old or echo genic thrombus low concns. of vesicles and vesicles targeted to tissues, cells or receptors.
 RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 15 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 2000:402087 CAPLUS
 DN 133:39968
 TI Separation of photosensitizer isomers and stereoisomers by laser-induced fluorescence capillary electrophoresis
 IN Dolphin, David; Peng, Xuejun; Sternbert, Ethan D.
 PA The University of British Columbia, Can.
 SO PCT Int. Appl., 50 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000034763	A2	20000615	WO 1999-US29352	19991210
	WO 2000034763	A3	20001012		
	W:				
	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
	RW:				
	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	CA 2353827	A1	20000615	CA 1999-2353827	19991210
	CA 2353827	C	20070626		
	EP 1137933	A2	20011004	EP 1999-965213	19991210
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	JP 2002532682	T	20021002	JP 2000-587170	19991210
	US 20020079221	A1	20020627	US 2001-26980	20011217
	US 6878253	B2	20050412		
PRAI	US 1998-111955P	P	19981211		
	US 1999-321893	A	19990528		
	WO 1999-US29352	W	19991210		
IT	361-09-1, Sodium cholate				
	RL: ARU (Analytical role, unclassified); ANST (Analytical study)				
	(chiral selector; separation of BPD-MA from its liposomal formulation)				
RN	361-09-1 CAPLUS				
CN	Cholan-24-oic acid, 3,7,12-trihydroxy-, sodium salt (1:1), (3 α ,5 β ,7 α ,12 α)- (CA INDEX NAME)				

Absolute stereochemistry.



● Na


AB A method for the separation of isomers and stereoisomers of photosensitizers by Laser-Induced Fluorescence Capillary Electrophoresis has been developed. The limits of detection are 2.06×10^{-6} M, and the relative standard deviation for the separation was 2.90 % to 4.64 %. Benzoporphyrin derivative mono and diacid

(BPD-MA, BPD-DA) enantiomers can be quant. determined in the range of 10^{-2} to 10^{-5} mg mL⁻¹. In comparison with HPLC, CE has better resolution and efficiency. This separation method was successfully applied to the BPD enantiomers obtained from a matrix of bovine serum and from liposomally formulated material as well as from studies with rat, dog and human microsomes.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 16 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
AN 1999:491468 CAPLUS
DN 131:303296
TI Biopharmaceutics of Boronated Radiosensitizers: Liposomal Formulation of MnBOPP (Manganese Chelate of 2,4-(α,β -Dihydroxyethyl) Deuterioporphyrin IX) and Comparative Toxicity in Mice
AU Zhou, Rong; Balasubramanian, Sathyamangalam V.; Kahl, Stephen B.; Straubinger, Robert M.
CS Department of Pharmaceutics, University at Buffalo State University of New York, Amherst, NY, 14260-1200, USA
SO Journal of Pharmaceutical Sciences (1999), 88(9), 912-917
CODEN: JPMSAE; ISSN: 0022-3549
PB American Chemical Society
DT Journal
LA English
IT 57-88-5, Cholesterol, biological studies
RL: BPR (Biological process); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)
(liposomal formulation of manganese chelate of 2,4-(α,β -dihydroxyethyl)deuterioporphyrin IX and comparative toxicity in mice)
RN 57-88-5 CAPLUS
CN Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)

Absolute stereochemistry.



AB Binary treatment modalities such as photodynamic therapy (PDT) and neutron capture therapy (NCT) combine low-toxicity electromagnetic irradiation with an appropriate radiation sensitizer to enhance selectivity for tumor targets. The porphyrin derivative tetrakis(carborane carboxylate ester of 2,4-(α,β -dihydroxyethyl) deuteroporphyrin IX (BOPP) shows tumor-selective uptake and is active in both treatment modalities. BOPP also chelates paramagnetic ions such as Mn^{2+} , and therefore its tissue accumulation and selectivity can be detected noninvasively by using magnetic resonance imaging. However, local and systemic toxicity appears elevated for the Mn^{2+} chelate (MnBOPP), but is poorly characterized. Here we have developed a liposomal formulation of MnBOPP and compared its toxicity with that of MnBOPP administered to mice in saline. The optimal liposome composition and maximal capacity to accommodate MnBOPP were investigated by differential scanning calorimetry and by encapsulation efficiency. MnBOPP was encapsulated quant. at up to 12 mol % (drug:lipid) in liposomes of varying composition, and remained incorporated during extended dialysis. Phase separation of drug- and lipid-rich domains was observed

above 12% drug. MnBOPP in buffered saline was lethal to animals at 90 $\mu\text{mol/kg}$, and caused severe necrosis at the injection site at dose levels of 60 $\mu\text{mol/kg}$ or greater. In contrast, MnBOPP formulated in liposomes was well tolerated at the highest tested dose of 135 $\mu\text{mol/kg}$, with the elimination of local toxicity.

RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 17 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN

AN 1998:728539 CAPLUS

DN 130:1847

TI Methods for treating viral infections with liposome-formulated photosensitizers

IN Ben-Hur, Ehud

PA New York Blood Center, Inc., USA

SO PCT Int. Appl., 22 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9849281	A1	19981105	WO 1998-US8479	19980428
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW			
	RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI,			

CM, GA, GN, ML, MR, NE, SN, TD, TG

US 6103706	A	20000815	US 1997-841042	19970429
AU 9871654	A	19981124	AU 1998-71654	19980428
US 6348453	B1	20020219	US 2000-543607	20000405
PRAI US 1997-841042	A	19970429		
WO 1998-US8479	W	19980428		

IT 57-88-5, Cholesterol, biological studies
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (liposome-formulated photosensitizers for treating viral
 infections)
 RN 57-88-5 CAPLUS
 CN Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)

Absolute stereochemistry.



AB The invention provides a method for treating a viral infection in a subject in need of such treatment which comprises administering to the subject a photosensitizer formulated in a liposome carrier, and exposing the subject to light at a wavelength 20-40 nm greater than the maximum absorption of the photosensitizer at a sufficient dose and duration to treat the viral infection in the subject. The invention also provides a method for treating a viral infection in a subject in need of such treatment comprising administering to the subject (i) a photosensitizer formulated in a liposome carrier and (ii) at least one quencher, and exposing the subject to light at a sufficient wavelength, dose, and duration to treat the viral infection in the subject.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 18 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 1998:484928 CAPLUS
 DN 129:113548
 OREF 129:23207a,23210a
 TI Pharmaceutical or cosmetic compositions containing homogeneously charged particulate vector
 IN Betbeder, Didier; Major, Michel
 PA Biovector Therapeutics S.A., Fr.
 SO PCT Int. Appl., 47 pp.
 CODEN: PIXXD2
 DT Patent
 LA French
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	WO 9829102	A1	19980709	WO 1997-FR2397	19971223
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,				

UA, UG, US, UZ, VN, YU, ZW
 RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI,
 FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM,
 GA, GN, ML, MR, NE, SN, TD, TG

FR 2757768	A1	19980703	FR 1996-16146	19961227
FR 2757768	B1	19990402		
CA 2276692	A1	19980709	CA 1997-2276692	19971223
AU 9856688	A	19980731	AU 1998-56688	19971223
EP 946153	A1	19991006	EP 1997-952990	19971223

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, FI


JP 2001508425	T	20010626	JP 1998-529682	19971223
---------------	---	----------	----------------	----------

PRAI FR 1996-16146 A 19961227
 WO 1997-FR2397 W 19971223

IT 57-88-5, Cholesterol, biological studies
 RL: BUU (Biological use, unclassified); THU (Therapeutic use); BIOL
 (Biological study); USES (Uses)
 (pharmaceutical or cosmetic compns. containing homogeneously charged
 particulate vector)

RN 57-88-5 CAPLUS
 CN Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)

Absolute stereochemistry.



AB The invention concerns a particulate carrier comprising a non-liquid hydrophilic nucleus; an amphiphilic lamella characterized in that the nucleus carries a global cationic, anionic or neutral charge and that the amphiphilic lamella carries a global charge of same polarity as that carried by the nucleus. The invention also concerns a pharmaceutical or cosmetic composition or a nutrient additive containing such a vector. Thus, maltodextrin (500 g) was treated with 7 g NaBH₄ followed by the reaction with NaOH, 30.25 mL epichlorohydrin and 382.3 g glycidyltrimethylammonium chloride. The resulting gel was diluted with water and neutralized with HOAc. Nanoparticle carriers were prepared by using the above polysaccharide and a phospholipid.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 19 OF 19 CAPLUS COPYRIGHT 2009 ACS on STN
 AN 1998:1394 CAPLUS
 DN 128:72421
 OREF 128:14083a,14086a
 TI Texaphyrin-lipophilic molecule-vesicle complexes, membrane incorporation of texaphyrins, and use in diagnosis and therapy
 IN Young, Stuart W.; Wright, Meredith; Sessler, Jonathan L.; Mody, Tarak D.; Magda, Darren
 PA Pharmacyclics, Inc., USA; Board of Regents, University of Texas System; Young, Stuart W.; Wright, Meredith; Sessler, Jonathan L.; Mody, Tarak D.; Magda, Darren

SO PCT Int. Appl., 67 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9746262	A2	19971211	WO 1997-US9501	19970604
	WO 9746262	A3	19980312		
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN			
	RW:	GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
	CA 2257225	A1	19971211	CA 1997-2257225	19970604
	AU 9732264	A	19980105	AU 1997-32264	19970604
	AU 727138	B2	20001207		
	CN 1225591	A	19990811	CN 1997-196446	19970604
	EP 954336	A2	19991110	EP 1997-927923	19970604
	EP 954336	B1	20040225		
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
	BR 9710685	A	20000111	BR 1997-10685	19970604
	NZ 333072	A	20000623	NZ 1997-333072	19970604
	JP 2000512279	T	20000919	JP 1998-500764	19970604
	IL 127315	A	20030529	IL 1997-127315	19970604
	AT 260121	T	20040315	AT 1997-927923	19970604
	US 20020006378	A1	20020117	US 1997-975090	19971120
	MX 9810198	A	20000531	MX 1998-10198	19981203
PRAI	US 1996-56917P	P	19960604		
	US 1996-657947	A	19960604		
	WO 1997-US9501	W	19970604		

OS MARPAT 128:72421

IT 57-88-5D, Cholesterol, texaphyrin conjugates, vesicle complexes


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)

(texaphyrin-lipophilic mol.-vesicle complexes, membrane incorporation of texaphyrins, and use in diagnosis and therapy)

RN 57-88-5 CAPLUS

CN Cholest-5-en-3-ol (3 β)- (CA INDEX NAME)

Absolute stereochemistry.



AB Compns. are provided having a texaphyrin-lipophilic mol. conjugate loaded into a biol. vesicle, as are methods for imaging, diagnosis and treatment using the loaded vesicle are provided. For example, liposomes

or red blood cells loaded with a paramagnetic texaphyrin-lipophilic mol. conjugate have utility as a blood pool contrast agent, facilitating the enhancement of normal tissues, magnetic resonance angiog., and marking areas of damaged endothelium by their egress through fenestrations or damaged portions of the blood vascular system. Liposomes or cells loaded with a photosensitive texaphyrin-lipophilic mol. conjugate can be photolyzed, allowing for a photodynamic therapy effect at the site of lysis. Availability of red blood cells loaded with a photosensitive texaphyrin-lipophilic mol. conjugate provides a method for delivering a photodynamic therapeutic agent to a desired site with a high concentration of oxygen. By presenting the agent in this way, it is expected that a patient will experience less toxicity.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> log

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y

(FILE 'HOME' ENTERED AT 17:16:10 ON 13 MAY 2009)

FILE 'CAPLUS' ENTERED AT 17:16:20 ON 13 MAY 2009

L1 0 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON 2006-591658/APPS
L2 1 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON US 20080269184/PN
SELECT L2 1 RN
L3 296106 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON (112-80-1/BI OR
1121-60-4/BI OR 1338-43-8/BI OR 143-02-2/BI OR 143-03-3/BI OR
143-07-7/BI OR 14982-53-7/BI OR 151-41-7/BI OR 313-04-2/BI OR
361-09-1/BI OR 40904-90-3/BI OR 4754-44-3/BI OR 516-95-0/BI OR
544-63-8/BI OR 57-10-3/BI OR 57-11-4/BI OR 57-88-5/BI OR
6156-78-1/BI OR 65028-70-8/BI OR 67-97-0/BI OR 691397-13-4/BI
OR 71794-64-4/BI OR 72924-08-4/BI OR 7789-46-0/BI OR 80-97-7/BI
OR 823808-59-9/BI OR 864444-61-1/BI OR 872-85-5/BI OR
9005-65-6/BI OR 9005-67-8/BI)

FILE 'REGISTRY' ENTERED AT 17:18:18 ON 13 MAY 2009

L4 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON 14982-53-7/RN
SET NOTICE 1 DISPLAY
D L4 SQIDE 1-
SET NOTICE LOGIN DISPLAY
L5 1 SEA FILE=REGISTRY SPE=ON ABB=ON PLU=ON L3 AND PORPHYRIN
D

FILE 'CAPLUS' ENTERED AT 17:19:22 ON 13 MAY 2009

L6 758 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L3 AND PORPHYRIN
L7 241 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L3 AND NIOSOME
L8 2 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L6 AND L7
D 1-2 BIB ABS
SET LINE 250
SET DETAIL OFF
E LIPOSOME+ALL/CT
SET LINE LOGIN
SET DETAIL LOGIN
L9 10368 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L3 AND ((LIPOSOME OR
"PHARMACEUTICAL LIPOSOMES") OR "LIPOSOMES")
L10 59 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L6 AND L9
L11 19 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L10 AND DRUG DELIVERY
D 1-19 BIB HITSTR ABS

COST IN U.S. DOLLARS

SINCE FILE TOTAL
ENTRY SESSION

FULL ESTIMATED COST	139.56	225.71
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-17.22	-17.22

STN INTERNATIONAL LOGOFF AT 17:35:15 ON 13 MAY 2009

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSPTAALF1626

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	DEC 01	ChemPort single article sales feature unavailable
NEWS	3	FEB 02	Simultaneous left and right truncation (SLART) added for CERAB, COMPUAB, ELCOM, and SOLIDSTATE
NEWS	4	FEB 02	GENBANK enhanced with SET PLURALS and SET SPELLING
NEWS	5	FEB 06	Patent sequence location (PSL) data added to USGENE
NEWS	6	FEB 10	COMPENDEX reloaded and enhanced
NEWS	7	FEB 11	WTEXTILES reloaded and enhanced
NEWS	8	FEB 19	New patent-examiner citations in 300,000 CA/CAPLUS patent records provide insights into related prior art
NEWS	9	FEB 19	Increase the precision of your patent queries -- use terms from the IPC Thesaurus, Version 2009.01
NEWS	10	FEB 23	Several formats for image display and print options discontinued in USPATFULL and USPAT2
NEWS	11	FEB 23	MEDLINE now offers more precise author group fields and 2009 MeSH terms
NEWS	12	FEB 23	TOXCENTER updates mirror those of MEDLINE - more precise author group fields and 2009 MeSH terms
NEWS	13	FEB 23	Three million new patent records blast AEROSPACE into STN patent clusters
NEWS	14	FEB 25	USGENE enhanced with patent family and legal status display data from INPADOCDB
NEWS	15	MAR 06	INPADOCDB and INPAFAMDB enhanced with new display formats
NEWS	16	MAR 11	EPFULL backfile enhanced with additional full-text applications and grants
NEWS	17	MAR 11	ESBIOBASE reloaded and enhanced
NEWS	18	MAR 20	CAS databases on STN enhanced with new super role for nanomaterial substances
NEWS	19	MAR 23	CA/CAPLUS enhanced with more than 250,000 patent equivalents from China
NEWS	20	MAR 30	IMSPATENTS reloaded and enhanced
NEWS	21	APR 03	CAS coverage of exemplified prophetic substances enhanced
NEWS	22	APR 07	STN is raising the limits on saved answers
NEWS	23	APR 24	CA/CAPLUS now has more comprehensive patent assignee

information

NEWS 24 APR 26 USPATFULL and USPAT2 enhanced with patent assignment/reassignment information

NEWS 25 APR 28 CAS patent authority coverage expanded

NEWS 26 APR 28 ENCOMPLIT/ENCOMPLIT2 search fields enhanced

NEWS 27 APR 28 Limits doubled for structure searching in CAS REGISTRY

NEWS 28 MAY 08 STN Express, Version 8.4, now available

NEWS 29 MAY 11 STN on the Web enhanced

NEWS 30 MAY 11 BEILSTEIN substance information now available on STN Easy

NEWS 31 MAY 14 DGENE, PCTGEN and USGENE enhanced with increased limits for exact sequence match searches and introduction of free HIT display format

NEWS 32 MAY 15 INPADOCDB and INPAFAMDB enhanced with Chinese legal status data

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN customer agreement. This agreement limits use to scientific research. Use for software development or design, implementation of commercial gateways, or use of CAS and STN data in the building of commercial products is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 16:02:10 ON 19 MAY 2009

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.22	0.22

FILE 'REGISTRY' ENTERED AT 16:02:20 ON 19 MAY 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 18 MAY 2009 HIGHEST RN 1147182-17-9
DICTIONARY FILE UPDATES: 18 MAY 2009 HIGHEST RN 1147182-17-9

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of

experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> file capl

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.48	0.70

FILE 'CAPLUS' ENTERED AT 16:02:28 ON 19 MAY 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 May 2009 VOL 150 ISS 21
FILE LAST UPDATED: 18 May 2009 (20090518/ED)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2009
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2009

Caplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate

=> s us 20080269184/pn

L1 1 US 20080269184/PN
(US20080269184/PN)

=> select l1

ENTER ANSWER NUMBER OR RANGE (1-):1

ENTER DISPLAY CODE (TI) OR ?:rn

E1 THROUGH E30 ASSIGNED

=> s e1-e30 and (niosome or liposome OR "Liposomes" OR "Pharmaceutical liposomes")
'E30' NOT FOUND

The E# entered is not currently defined.

=> s e1-e30

'E30' NOT FOUND

The E# entered is not currently defined.

=> s e1-e29

'E29' NOT FOUND

The E# entered is not currently defined.

=> sel

SE1 IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system.
For a list of commands available to you in the current file, enter
"HELP COMMANDS" at an arrow prompt (=>).

=> s e1

L2 12656 LIPOSOME/CT

=> s e2

L3 8315 LIPOSOMES/CT

=> select l1

ENTER ANSWER NUMBER OR RANGE (1-):1

ENTER DISPLAY CODE (TI) OR ?:rn

E1 THROUGH E30 ASSIGNED

=> s e1-e30

58767 112-80-1/BI

4779 1121-60-4/BI

6325 1338-43-8/BI

138 143-02-2/BI

87 143-03-3/BI

21482 143-07-7/BI

297 14982-53-7/BI

1357 151-41-7/BI

1167 313-04-2/BI

2376 361-09-1/BI

60 40904-90-3/BI

137 4754-44-3/BI

396 516-95-0/BI

26179 544-63-8/BI

51564 57-10-3/BI

63460 57-11-4/BI

141436 57-88-5/BI

488 6156-78-1/BI

11 65028-70-8/BI

7081 67-97-0/BI

5430 691397-13-4/BI

43 71794-64-4/BI

54 72924-08-4/BI

766 7789-46-0/BI

2644 80-97-7/BI

3 823808-59-9/BI

1 864444-61-1/BI

3773 872-85-5/BI

17843 9005-65-6/BI

4772 9005-67-8/BI

L4 296567 (112-80-1/BI OR 1121-60-4/BI OR 1338-43-8/BI OR 143-02-2/BI OR
143-03-3/BI OR 143-07-7/BI OR 14982-53-7/BI OR 151-41-7/BI OR
313-04-2/BI OR 361-09-1/BI OR 40904-90-3/BI OR 4754-44-3/BI OR
516-95-0/BI OR 544-63-8/BI OR 57-10-3/BI OR 57-11-4/BI OR 57-88-
5/BI OR 6156-78-1/BI OR 65028-70-8/BI OR 67-97-0/BI OR 691397-13
-4/BI OR 71794-64-4/BI OR 72924-08-4/BI OR 7789-46-0/BI OR 80-97
-7/BI OR 823808-59-9/BI OR 864444-61-1/BI OR 872-85-5/BI OR 9005
-65-6/BI OR 9005-67-8/BI)

=> s l4 and (niosome or liposome OR "Liposomes" OR "Pharmaceutical liposomes")

360 NIOSOME

481 NIOSOMES

502 NIOSOME

(NIOSOME OR NIOSOMES)

40392 LIPOSOME

```

52258 LIPOSOMES
60040 LIPOSOME
      (LIPOSOME OR LIPOSOMES)
52258 "LIPOSOMES"
342378 "PHARMACEUTICAL"
93571 "PHARMACEUTICALS"
398332 "PHARMACEUTICAL"
      ("PHARMACEUTICAL" OR "PHARMACEUTICALS")
52258 "LIPOSOMES"
5005 "PHARMACEUTICAL LIPOSOMES"
      ("PHARMACEUTICAL"(W)"LIPOSOMES")
L5      10513 L4 AND (NIOSOME OR LIPOSOME OR "LIPOSOMES" OR "PHARMACEUTICAL
      LIPOSOMES")

```

```

=> s l4 and niosome
      360 NIOSOME
      481 NIOSOMES
      502 NIOSOME
      (NIOSOME OR NIOSOMES)
L6      241 L4 AND NIOSOME

```

```

=> s l4 and (niosome/ab or niosome/ti)
      127 NIOSOME/AB
      346 NIOSOMES/AB
      391 NIOSOME/AB
      ((NIOSOME OR NIOSOMES)/AB)
      59 NIOSOME/TI
      177 NIOSOMES/TI
      235 NIOSOME/TI
      ((NIOSOME OR NIOSOMES)/TI)
L7      210 L4 AND (NIOSOME/AB OR NIOSOME/TI)

```

```

=> s l4 and or niosome/ti
MISSING TERM 'AND OR'
The search profile that was entered contains a logical
operator followed immediately by another operator.

```

```

=> s l4 and niosome/ti
      59 NIOSOME/TI
      177 NIOSOMES/TI
      235 NIOSOME/TI
      ((NIOSOME OR NIOSOMES)/TI)
L8      138 L4 AND NIOSOME/TI

```

```

=> s l5 and (porphyrin or metalloporphyrin)
      40679 PORPHYRIN
      27805 PORPHYRINS
      47740 PORPHYRIN
      (PORPHYRIN OR PORPHYRINS)
      4470 METALLOPORPHYRIN
      7135 METALLOPORPHYRINS
      8719 METALLOPORPHYRIN
      (METALLOPORPHYRIN OR METALLOPORPHYRINS)
L9      63 L5 AND (PORPHYRIN OR METALLOPORPHYRIN)

```

```

=> s l9 and niosome
      360 NIOSOME
      481 NIOSOMES
      502 NIOSOME
      (NIOSOME OR NIOSOMES)
L10     2 L9 AND NIOSOME

```


=> d 1-2 ibib abs

L10 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:1004559 CAPLUS

DOCUMENT NUMBER: 143:292573

TITLE: Niosome having metal porphyrin
complex embedded therein, process for producing the
same and drug with the use thereof

INVENTOR(S): Yuasa, Makoto; Oyaizu, Kenichi; Yamaguchi, Aritomo;
Hanyuu, Yukihiro; Kasahara, Kazunori; Komuro, Masayasu

PATENT ASSIGNEE(S): Japan

SOURCE: PCT Int. Appl., 40 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
WO 2005084665	A1	20050915	WO 2004-JP2750	20040304
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1731150	A1	20061213	EP 2004-717289	20040304
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
CN 1942184	A	20070404	CN 2004-80042914	20040304
KR 2007008623	A	20070117	KR 2006-720709	20061002
US 20080269184	A1	20081030	US 2007-591658	20070815
PRIORITY APPLN. INFO.:			WO 2004-JP2750	W 20040304

OTHER SOURCE(S): MARPAT 143:292573

AB Disclosed is a niosome having a metal porphyrin
complex embedded therein which contains a cationized metal
porphyrin complex and a niosome-forming substance. This
niosome having a metal porphyrin complex embedded
therein has an SOD activity and can target super oxide anion radical
(O₂⁻.) and surely decrease it. Because of being in the form of a
niosome, it can be delivered to, for example, a cancer cell in
vivo. Therefore, it can decrease O₂⁻ in a cancer cell and exert an
excellent effect of treating cancer. Moreover, it shows a selective
effect and, therefore, is usable as a novel anticancer agent with no side
effect. In addition, it can be hold in the blood, which makes it favorable
as an antioxidant. Owing to this characteristic, it can protect the
living body from in vivo disorders caused by active oxygen. For example,
iron[5,10,15,20-tetrakis(2-methylpyridyl)porphyrin] was prepared,
and mixed with stearic acid metal salt to form an ion complex of the
porphyrin. Then, the ion complex was mixed with tween-61 and
cholesterol to form a niosome to exam for its antitumor activity
and antioxidant activity in vitro.

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2005:674689 CAPLUS
 DOCUMENT NUMBER: 144:156398
 TITLE: Novel functional nano-size nonionic surfactant particles on which cationic metalloporphyrins are absorbed; preparation, characterization, and antioxidant properties
 AUTHOR(S): Yuasa, Makoto; Oyaizu, Kenichi; Hanyuu, Yukihiro; Kasahara, Kazunori; Yamaguchi, Aritomo
 CORPORATE SOURCE: Department of Pure and Applied Chemistry, Faculty of Science and Technology, Tokyo University of Science, Noda, 278-8510, Japan
 SOURCE: Journal of Oleo Science (2005), 54(8), 465-471
 CODEN: JOSOAP; ISSN: 1345-8957
 PUBLISHER: Japan Oil Chemists' Society
 DOCUMENT TYPE: Journal
 LANGUAGE: Japanese
 AB Cationic manganese(III) 5,10,15,20-tetrakis(N-methylpyridinium-2-yl) porphyrin (MnT2MPyP) and manganese(III) 5,10,15,20-tetrakis(N-methylpyridinium-4-yl) porphyrin (MnT4MPyP) complexes were synthesized as superoxide dismutase (SOD) mimics which were introduced into niosomes to examine this effects on the capacity of drug delivery system (DDS) to maintain and perpetuate blood circulation. All the niosomes were prepared from polyoxyethylene sorbitan monostearate (Tween 61) by the conventional sonication method. SOD activity was measured by the stopped-flow anal. and the cytochrome c method. Sodium stearate-linked MnT2MPyP was the most effective catalyst along with SOD activity for decomposing O₂⁻ at a second-order rate constant of 2.0×10⁷ M⁻¹ s⁻¹ in Tween 61 niosomes. Rate consts. of metalloporphyrin-embedded niosomes for reaction with hydrogen peroxide (H₂O₂) and half-life times in H₂O₂ were also determined Metalloporphyrin-embedded niosomes were found to have greater half-life times compared to metalloporphyrin without niosomes. The present findings clearly indicate that metalloporphyrin-embedded niosomes are highly effective for bringing about O₂⁻ decomposition and should thus find application as DDS in antioxidant drugs.

=> s 19 and (nonionic or non-ionic) surfactant
 MISSING OPERATOR ON-IONIC) SURFACTANT
 The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> s 19 and (nonionic or non-ionic) (A) surfactant
 79673 NONIONIC
 543 NONIONICS
 79854 NONIONIC
 (NONIONIC OR NONIONICS)
 1053499 NON
 38 NONS
 1053528 NON
 (NON OR NONS)
 307262 IONIC
 545 IONICS
 307545 IONIC
 (IONIC OR IONICS)
 9415 NON-IONIC
 (NON(W) IONIC)
 215520 SURFACTANT
 192503 SURFACTANTS
 274903 SURFACTANT
 (SURFACTANT OR SURFACTANTS)

52477 (NONIONIC OR NON-IONIC) (A) SURFACTANT
L11 2 L9 AND (NONIONIC OR NON-IONIC) (A) SURFACTANT

=> s l11 not l10
L12 0 L11 NOT L10

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	120.46	121.16
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-1.64	-1.64

FILE 'STNGUIDE' ENTERED AT 16:07:34 ON 19 MAY 2009
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: May 15, 2009 (20090515/UP).

=> log
ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF
LOGOFF? (Y)/N/HOLD:y

(FILE 'HOME' ENTERED AT 16:02:10 ON 19 MAY 2009)

FILE 'REGISTRY' ENTERED AT 16:02:20 ON 19 MAY 2009

FILE 'CAPLUS' ENTERED AT 16:02:28 ON 19 MAY 2009

L1 1 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON US 20080269184/PN
SELECT L1 1 RN
SET LINE 250
SET DETAIL OFF
E LIPOSOME+ALL/CT
SET LINE LOGIN
SET DETAIL LOGIN

L2 12656 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON LIPOSOME/CT
L3 8315 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON LIPOSOMES/CT
SELECT L1 1 RN

L4 296567 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON (112-80-1/BI OR
1121-60-4/BI OR 1338-43-8/BI OR 143-02-2/BI OR 143-03-3/BI OR
143-07-7/BI OR 14982-53-7/BI OR 151-41-7/BI OR 313-04-2/BI OR
361-09-1/BI OR 40904-90-3/BI OR 4754-44-3/BI OR 516-95-0/BI OR
544-63-8/BI OR 57-10-3/BI OR 57-11-4/BI OR 57-88-5/BI OR
6156-78-1/BI OR 65028-70-8/BI OR 67-97-0/BI OR 691397-13-4/BI
OR 71794-64-4/BI OR 72924-08-4/BI OR 7789-46-0/BI OR 80-97-7/BI
OR 823808-59-9/BI OR 864444-61-1/BI OR 872-85-5/BI OR
9005-65-6/BI OR 9005-67-8/BI)

L5 10513 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L4 AND (NIOsome OR
LIPOSOME OR "LIPOSOMES" OR "PHARMACEUTICAL LIPOSOMES")

L6 241 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L4 AND NIOSOME
L7 210 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L4 AND (NIOsome/AB OR
NIOsome/TI)

L8 138 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L4 AND NIOSOME/TI
SET LINE 250
SET DETAIL OFF
E PORPHYRIN+ALL/CT
SET LINE LOGIN
SET DETAIL LOGIN

L9 63 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L5 AND (PORPHYRIN OR
 METALLOPORPHYRIN)
 L10 2 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L9 AND NIOSOME
 D 1-2 IBIB ABS
 L11 2 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L9 AND (NONIONIC OR
 NON-IONIC) (A) SURFACTANT
 L12 0 SEA FILE=CAPLUS SPE=ON ABB=ON PLU=ON L11 NOT L10

FILE 'STNGUIDE' ENTERED AT 16:07:34 ON 19 MAY 2009

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.70	121.86
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-1.64

STN INTERNATIONAL LOGOFF AT 16:13:20 ON 19 MAY 2009